

**General information**



Wellbore name	24/9-5
Type	EXPLORATION
Purpose	WILDCAT
Status	P&A
Factmaps in new window	<a href="#">link to map</a>
Main area	NORTH SEA
Field	<a href="#">VOLUND</a>
Discovery	<a href="#">24/9-5 Volund</a>
Well name	24/9-5
Seismic location	FI 89 3D INLINE 1070 & CROSSLINE 2792
Production licence	<a href="#">150</a>
Drilling operator	Fina Production Licenses AS
Drill permit	778-L
Drilling facility	<a href="#">WEST DELTA</a>
Drilling days	51
Entered date	07.12.1993
Completed date	26.01.1994
Release date	26.01.1996
Publication date	15.02.2006
Purpose - planned	WILDCAT
Reentry	NO
Content	OIL
Discovery wellbore	YES
1st level with HC, age	EOCENE
1st level with HC, formation	INTRA BALDER FM SS
Kelly bushing elevation [m]	29.0
Water depth [m]	122.0
Total depth (MD) [m RKB]	2860.0
Maximum inclination [°]	5.9
Bottom hole temperature [°C]	88
Oldest penetrated age	LATE CRETACEOUS
Oldest penetrated formation	JORSALFARE FM
Geodetic datum	ED50
NS degrees	59° 29' 6.52" N
EW degrees	1° 55' 10.82" E
NS UTM [m]	6594725.25
EW UTM [m]	438806.71
UTM zone	31
NPDID wellbore	2244

## Wellbore history

### General

Wildcat well 24/9-5 was drilled in a location SW of the Heimdal field, ca 5 km from the UK border. It was programmed to test two primary sandstone prospects in the Late Paleocene. The lower objective was the Hermod Formation and the upper objective was sandstones within the Balder Formation. The Hermod Formation prospect was a mapped isochron thick with a small area of structural closure at Top Sele Formation. The Balder Formation prospect was mapped as an isochron thick with an associated seismic amplitude anomaly and was primarily regarded as a stratigraphic trap.

### Operations and results

Wildcat well 24/9-5 was spudded with the semi-submersible installation West Delta on 7 December 1993 and drilled to TD at 2860 m in the Late Cretaceous Jorsalfare Formation. No significant problem was reported from the operations. The well was drilled with seawater down to 581 m, with KCI/PAC/PHPA from 581 m to 1941 m, and with KCI/PAC/PHPA/glycol from 1941 m to TD.

The Balder Formation was encountered at 1964 m. A gross oil bearing Intra Balder Formation sandstone interval of 7.1 m was encountered at 2011 m. Net sandstone was 6.3 m of which 5.9 m was in a single massive unit. Petrophysical analysis gave porosities of 35 - 40% in the massive sandstone unit with an average SW of 22.2%. Base of the reservoir was encountered at 2018 m, which was found to be an oil-down-to-water contact. Samples from the FMT tool indicated the reservoir fluid to consist of 32-34 deg API oil with a GOR of 91 Sm<sup>3</sup>/Sm<sup>3</sup>. Pressure gradient analysis gave a free water level at 2031.5 m (2028.0 m TVD). Geochemical analysis of the oil sample indicated minor biodegradation had taken place. The Sele Formation was encountered at 2031 m with 47 m net sandstone in two Hermod Formation sequences. The upper sequence was encountered at 2049 m and the lower at 2098 m. These sandstones proved to be water bearing with no shows. A thin sandstone at 2045.5 - 2047 m, just above the upper Hermod interval, gave the highest known water below the Balder Formation pay interval. An additional zone of interest was encountered in the interval 1775-1790 m where a number of siltstone/argillaceous sandstone stringers were identified. Minor gas shows were observed and rare traces of sandstone gave poor (residual?) oil shows. Log analysis showed the presence of hydrocarbons in thin argillaceous stringers particularly over the interval 1785-1787.5 m where neutron density logs (acquired through casing) indicated the presence of gas (Average SW = 40%).

Four conventional cores were cut. Three were cut in the Late Palaeocene Balder Formation with 7 m Intra Balder Formation sandstone. The fourth was cut in the Hermod Formation Sandstone within the Sele Formation. Three FMT samples were taken in the well, one at 2013.5 m (oil) and two at 2017.5 m (mud filtrate)

Although the hydrocarbon-bearing interval in the Balder Formation was as prognosis it was recognised that the well was not in an optimum location and it was decided to drill an immediate appraisal well. Well 24/9-5 was therefore not tested.

The well was permanently abandoned on 26 January 1994 as an oil discovery.

### Testing

No drill stem test was performed in the well.

### Cuttings at the Norwegian Offshore Directorate

Cutting sample, top depth [m]	Cutting samples, bottom depth [m]
590.00	2860.00

Cuttings available for sampling?	YES
----------------------------------	-----

### Cores at the Norwegian Offshore Directorate

Core sample number	Core sample - top depth	Core sample - bottom depth	Core sample depth - uom
1	2000.0	2001.8	[m ]
2	2002.5	2019.9	[m ]
3	2020.3	2033.1	[m ]
4	2048.3	2052.8	[m ]

Total core sample length [m]	36.5
Cores available for sampling?	YES

### Core photos



2000-2002m



2002-2007m



2007-2012m



2012-2017m



2017-2020m



2020-2025m



2025-2030m



2030-2033m



2048-2052m

### Lithostratigraphy

Top depth [mMD RKB]	Lithostrat. unit
151	<a href="#">NORDLAND GP</a>
360	<a href="#">UTSIRA FM</a>

550	<a href="#">HORDALAND GP</a>
555	<a href="#">SKADE FM</a>
695	<a href="#">NO FORMAL NAME</a>
1203	<a href="#">GRID FM</a>
1255	<a href="#">NO FORMAL NAME</a>
1294	<a href="#">GRID FM</a>
1309	<a href="#">NO FORMAL NAME</a>
1964	<a href="#">ROGALAND GP</a>
1964	<a href="#">BALDER FM</a>
2011	<a href="#">INTRA BALDER FM SS</a>
2018	<a href="#">BALDER FM</a>
2031	<a href="#">SELE FM</a>
2049	<a href="#">HERMOD FM</a>
2082	<a href="#">SELE FM</a>
2098	<a href="#">HERMOD FM</a>
2112	<a href="#">SELE FM</a>
2116	<a href="#">LISTA FM</a>
2161	<a href="#">HEIMDAL FM</a>
2624	<a href="#">LISTA FM</a>
2728	<a href="#">TY FM</a>
2828	<a href="#">SHETLAND GP</a>
2828	<a href="#">JORSALFARE FM</a>

### Composite logs

Document name	Document format	Document size [MB]
<a href="#">2244</a>	pdf	0.36

### Geochemical information

Document name	Document format	Document size [MB]
<a href="#">2244_1</a>	pdf	1.88

### Documents - reported by the production licence (period for duty of secrecy expired)



Document name	Document format	Document size [MB]
<a href="#">2244_24_9_5_COMPLETION_REPORT_AND_LOG</a>	pdf	17.34

## Logs

Log type	Log top depth [m]	Log bottom depth [m]
ACBL	129	574
ACBL	450	1344
BRIDGEPLUG BST CCL	1700	1806
CBL VDL GR	1010	1932
CDL CN GR CAL	1750	2858
DIFL ACL SP GR CAL	129	1362
DIFL ACL SP GR CAL	1344	1881
DIFL ACL SP GR CAL	1932	2859
DLL MLL SL CAL SP	1930	2350
FMT VPC GR	2011	2320
FMT VPC GR	2012	2012
FMT VPC GR	2012	2111
FMT VPC GR	2017	2017
HDIP GR	1932	2350
MWD LWD - GR DPR	159	2834
SWC PFC	1950	2657
VSP	500	2850

## Casing and leak-off tests

Casing type	Casing diam. [inch]	Casing depth [m]	Hole diam. [inch]	Hole depth [m]	LOT/FIT mud eqv. [g/cm3]	Formation test type
CONDUCTOR	30	235.0	36	240.0	0.00	LOT
SURF.COND.	20	576.0	26	581.0	0.00	LOT
INTERM.	13 3/8	1346.0	17 1/2	1377.0	0.00	LOT
INTERM.	9 5/8	1935.0	12 1/2	1941.0	0.00	LOT
OPEN HOLE		2860.0	8 1/2	2860.0	0.00	LOT

## Drilling mud



Depth MD [m]	Mud weight [g/cm <sup>3</sup> ]	Visc. [mPa.s]	Yield point [Pa]	Mud type	Date measured
240	1.20	15.0		WATER BASED	
580	1.25	100.0		WATER BASED	
1124	1.17	18.0		WATER BASED	
1377	1.25	23.0		WATER BASED	
1377	1.22	20.0		WATER BASED	
2203	1.25	23.0		DUMMY	
2860	1.25	21.0		DUMMY	

### Pressure plots

The pore pressure data is sourced from well logs if no other source is specified. In some wells where pore pressure logs do not exist, information from Drill stem tests and kicks have been used. The data has been reported to the NPD, and further processed and quality controlled by IHS Markit.

Document name	Document format	Document size [MB]
<a href="#">2244_Formation_pressure_(Formasjonstrykk)</a>	pdf	0.21

